


Measuring Guide for Compression Stockings

An important measuring tip: If swelling is an issue, measure your legs first thing in the morning, after you get out of bed. This is when swelling is at a minimum and the size of your legs should remain once the swelling is eliminated.



1 Ankle circumference at narrowest part of ankle

2 Calf circumference at fullest part of the calf

3 Calf length from the floor to the right-angle bend of the knee

4 Widest circumference of the thigh

5 Leg length from the floor up to the gluteal fold

6 Circumference at the hip

Knee High

Select One: Open Toe or Closed Toe

Left Right

1. Ankle: _____
2. Calf: _____
3. Calf Length: _____

All:

Select One: Open Toe or Closed Toe

Left Right

1. Ankle: _____
2. Calf: _____
3. Calf Length: _____
4. Thigh: _____
5. Thigh Length: _____
6. Hip: _____

Thigh High

Select One: Open Toe or Closed Toe

Left Right

1. Ankle: _____
2. Calf: _____
4. Thigh: _____
5. Thigh Length: _____

Diagnosis: (Required)

of Refills: (max of one pair per month—11 refills)

Pantyhose

Select One: Open Toe or Closed Toe

Left Right

1. Ankle: _____
2. Calf: _____
4. Thigh: _____
5. Length: _____
6. Hips: _____

Compression Strength:

18-30 mmHg | 30-40 mmHg | 40-60 mmHG

Patient Info:

Customer Name: _____

Physical Address: _____

Date of Birth: _____

Phone#: _____

Signature of Physician, Nurse Practitioner, Physicians Assistant

Printed Name of Physician, Nurse Practitioner, Physicians Assistant

NPI#: _____

Date: _____

FAX FORM AND MEDICAL RECORDS TO 207-622-3264